

Scalar Curvature Rigidity 2

Thm 1 (Llarull '98)

(M^n, g) closed connected spin, $n \geq 2$

a) $\text{scal}_g \geq n(n-1) = \text{scal}_{g_{S^n}}$

b) $f: (M, g) \rightarrow (S^n, g_0)$ smooth, $\deg(f) \neq 0$ s.t.

• f 1-Lipschitz, $n = 2$

• f area-nonincreasing, $n \geq 3$

$\Rightarrow f$ Riem. isometry

- Goette-Sennemann '02: replace S^n by mfd. with positive Riem. curvature operator and non-vanishing Euler characteristic
- Lott '21: Goette-Sennemann for compact mfd. with boundary
- Brendle '24: Convex polytopes
- Bär-Brendle-Hanke-Wang '24: S^n with antipodal points removed

Thm 2 (Cecchini-Hanke-Schick '22, C-H-S-S '25)

M^n smooth closed connected spin, $n \geq 2$

a) $g \in W^{1,p}(T^*M \otimes T^*M)$, $p > n$, admissible Riem. metric s.t.

$$\text{scal}_g^{\text{dist.}} \geq n(n-1)$$

b) $f: (M, g) \rightarrow (S^n, g_0)$ Lipschitz, $\deg(f) \neq 0$ s.t.

- f 1-Lipschitz, $n=2$
- f area-nonincreasing a.e., $n \geq 3$

$\Rightarrow f: (M, d_g) \rightarrow (S^n, d_0)$ metric isometry

Remark Lee-Tam '22: Thm 2 in the case f is 1-Lipschitz $\forall n \geq 2$

Def M closed mfd.

- $0 < \nu \leq 1$, $g_r \in C^\infty((0, \nu), C^\infty(T^*M \otimes T^*M))$
family of Riem. metrics

- $g_0 \in C^2(T^*M \otimes T^*M)$

s.t. $g_r \xrightarrow{r \rightarrow 0} g_0$ in C^2 and

$$|r \partial_r g_r| \rightarrow 0, \quad |r^2 \partial_r^2 g_r| \rightarrow 0 \quad \text{as } r \rightarrow 0$$

Then $G_r := dr^2 + r^2 g_r$ on $(0, \nu) \times M$ is called generalized cone metric.

(W, G) is a smooth Riem. mfd. with cone-like singularities, if

- $\exists \underline{K} \subset W$ compact
- $\exists 0 < \nu \leq 1$ and Riem. isometry

$$\nu: (W \setminus K, G|_{W \setminus K}) \rightarrow ((0, \nu) \times \partial K, G_r)$$

↑
generalized
cone metric



Thm 3 (C-H-S-S '25)

(W^{n+1}, G) Riem. spin mfd. with cone-like sing.

$n \geq 3$ odd


- $\text{scal}_G \geq (n+1)n$
- $f: W \rightarrow S^{n+1}$ Lipschitz, area-nonincreasing a.e.
 $\deg(f) \neq 0$

$\Rightarrow f$ Riem. isometry onto a punctured sphere

Proof of Thm 2 part 2

local isometry $\leadsto$ metric isometry for n even

to show: f either or. pres. or or. rev. a.e.



$\rightarrow$ l.l. local isometry a.e. but

$$2 = d(-1, 1) \neq d(1-1, 1-1) = 0$$

$$0 \neq u \in \ker(D_E^+)$$

$$\Sigma M = \Sigma^+ M \oplus \Sigma^- M$$

From SL-formula

$$\bullet (e_i \cdot e_j \otimes f_* e_i \cdot f_* e_j) \cdot u = u \quad (1)$$

$$\bullet \nabla^{S \otimes E} u = 0 \quad \text{a.e.} \quad \leadsto |u| = \text{const. a.e.} \quad (2)$$

let $x \in M$ s.t. $d_x f$ exists

let $\omega_{E, n}, \omega_{F, n}$ be the volume forms

$$i^k e_1 \wedge \dots \wedge e_n$$

$$\Rightarrow (\omega_{\mathbb{C}, M} \otimes \omega_{\mathbb{C}, S^n}) \cdot u = u \quad (3)$$

$$f \text{ local isometry a.e.} \Rightarrow \omega_{\mathbb{C}, S^n} = \pm i^k f_* e_1 \wedge \dots \wedge f_* e_n$$

$\uparrow$
 $+$ if f or. pres.
 $-$ if f or. rev.

$$\Rightarrow \pm u \stackrel{(3)}{=} i^{2k} (e_1 \wedge \dots \wedge e_n \otimes f_* e_1 \wedge \dots \wedge f_* e_n) \cdot u$$

$$\stackrel{(1)}{=} (-1)^k u$$

$$\Rightarrow f \text{ orient} \begin{cases} \text{pres.} & \text{if } k \text{ even} \\ \text{rev.} & \text{if } k \text{ odd} \end{cases}$$

$$\stackrel{(2)}{\Rightarrow} \text{holds a.e.}$$

$$\Rightarrow f \text{ metric isometry}$$


Reshetnyak

n odd

Consider the spherical susp.

$$SM := M \times (0, \pi) \quad Sg := dr^2 + \sin^2(r)g$$

$$Sf : (SM, Sg) \longrightarrow (SS^n, Sg_n) = (S^{n+1} \setminus \{N, S\}, g_{S^{n+1}})$$



$$\text{scal}_{S_g} \geq (n+1)n = \text{scal}_{g_{S^{n+1}}}$$

But f area-nonincr. does not imply that

$$Sf = 1$$

Solution: first show f 1-Lipschitz using spectral flow

Then generalize even dim. proof to $SM \xrightarrow{Sf} S^{n+1}$

Caveat: Only have SL-formula for $p > n+1$ on SM

Spectral flow

"Def" Given a "continuous" path of self-adj. Fredholm op. D_t , $t \in [a, b]$, the spectral flow is

$sf(D_t) := \# \{ \text{eigenvalues passing from } - \text{ to } + \}$

$- \# \{ - \text{ to } + \} + \# \{ + \text{ to } - \}$

Thm (Getzler '93)

M^{2k+1} closed connected spin, D Dirac op. on $\Sigma M \otimes \mathbb{C}^N$
for $N \in \mathbb{N}$

Let $U: M \rightarrow U(N)$ smooth and

$$D_t := (1-t)D + t U^{-1} D U \quad , \quad t \in [0,1]$$

$$\leadsto \text{sf}(D_t) = - \frac{1}{(-2\pi i)^{k+1}} \int_M \hat{A}(TM) \wedge \text{Ch}(U)$$

$$\text{If } M = S^{2k+1} \quad \text{sf}(D_t) = \deg(U)$$

Def For $u \in C^\infty(\Sigma M)$ set $X \in TM$

$$\nabla_X^t u = \nabla_X^{\Sigma M} u + t X \cdot u$$

∇^t -parallel spinors are called Killing spinors

Example: $n=2k, n=2k+1$

For $t = \pm \frac{1}{2}$, ΣS^n can be trivialized by $\nabla^{\pm \frac{1}{2}}$ -parallel spinors i.e.

$$\underbrace{\exists \Sigma S^n \xrightarrow{V_\pm} S^n \times \mathbb{C}^{2^k}}_{(*)}$$

$f: M \rightarrow S^n$, consider $E = f^*(S^n \times \mathbb{C}^{2^k}) = M \times \mathbb{C}^{2^k}$
with trivial connection

$\leadsto D_E$ twisted Dirac op.

(*) induce trivializ. $U_{\pm} : E \xrightarrow{\sim} M \times \mathbb{C}^{2k}$

$$\text{Set } U = U_+ U_-^{-1} : M \rightarrow U(\mathbb{C}^{2k})$$

Li-Su-Wang

$$\Rightarrow \text{sf}(D_{\epsilon}) = \deg(f) \neq 0$$

$$\Rightarrow \text{for some } t \in [0, 1] \quad \exists u \in \ker(D_t)$$

Abstract cone operators

Consider $SM = (0, \pi) \times M$, $g = dr^2 + \sin^2(r)g$

There are bundle isometries

$$\Sigma^{\pm} SM|_{M \times \{x\}} \cong \Sigma M$$

$$\text{let } U = (0, 1) \times M$$

Using parallel transport of sections in $C^{\infty}(\Sigma M)$
along geodesic lines $(0, 1) \times \{x\} \subset SM$, $\Sigma^{\pm} SM$

we get isometries



_____ (0, 1)

$$L^2(U, \Sigma^\pm SM, d\text{vol}_{S_g}) \xrightarrow[\sim]{\phi_\pm}$$

$$\xrightarrow{\sim} L^2((0,1), L^2(M, \Sigma M, \sin^n(r) d\text{vol}_g))$$

s.t.

$$\phi_- \circ D^+ \circ \phi_-^{-1} = \partial_r + \frac{n}{2} \cot(r) + \frac{1}{\sin(r)} D_M$$

compose $\phi_\pm$ with mult. with $\sin^{n/2}(r)$

$$\left[(u, v)_{L^2} = \int_0^1 \int_M \underbrace{\langle u, v \rangle}_{\text{mult. with } \sin^{n/2}(r)} \sin^n(r) d\text{vol}_g dr \right]$$

$$\sim L^2(U, \Sigma^\pm SM, d\text{vol}_{S_g})$$

$$\xrightarrow[\sim]{\tilde{\phi}_\pm} L^2((0,1), \underbrace{L^2(M, \Sigma M; d\text{vol}_g)}_{=: L})$$

and

$$\begin{aligned} \tilde{\phi}_- \circ D^+ \circ \tilde{\phi}_+^{-1} &= \partial_r + \frac{1}{\sin(r)} D_M \quad \checkmark \text{ Dirac op. on } M \\ &= \partial_r + \frac{1}{r} D_M + \underbrace{\left(\frac{1}{\sin(r)} - \frac{1}{r} \right) D_M}_{\text{pert. term}} \end{aligned}$$

→ construct parametrix for $\bar{D}^+$

$P: L^2(\Sigma^+ SM) \longrightarrow \text{dom}(\bar{D}^+)$ bounded s.t.

$$P \bar{D}^+ = \text{id} + L \quad \bar{D}^+ P = \text{id} + R$$

compact operators

existence of parametrix for $\bar{D}^+$

$\Leftrightarrow \bar{D}^+$ Fredholm

deform. argument

$\Rightarrow$

$\exists$ harm. spinor

+ index formula

of Chou '85

Furthermore, integrated SL-formula still holds
for mfd's. with cone-like singularities