

Invariant theory of Hamiltonian mechanics and related numerical analysis

Oscar Cosserat

Göttingen Mathematisches Institut

IIITD, September 2025

Geometric mechanics



Isaac
**New-
ton**,
1643-
1727



Émilie
**du
Châtelet**,
1706-
1749



Joseph-
Louis **de
La-
grange**,
1736-
1813



Siméon
Denis
Poisson,
1781-
1840



Marius
Sophus
Lie,
1842-
1899



Vladimir
I.
Arnol'd,
1937-
2010



André
**Lich-
nerow-
icz**,
1915-
1998

and Carl Gustav Jacob **Jacobi** (1804 - 1851), Sir William Rowan **Hamilton** (1805 –1865), Sofia Vassilievna **Kovalevskaja** (1850-1891), Henri **Poincaré** (1854-1912), Jean-Marie **Souriau** (1922 - 2012), Alan David **Weinstein** (1943-)...

Hamiltonian dynamics on Poisson manifolds

- Generalities on Poisson structures

- Hamiltonian systems

- A first approach to the numerical analysis on Poisson manifolds

Symplectic groupoids

- Basic notions on groupoids

- Basics on symplectic groupoids

- Lagrangian bisections

- Birealisations

- Hamilton-Jacobi equation

Hamiltonian Poisson integrators

- Butcher series

- Explicit construction through birealisations

- Numerical tests around a singularity: a Lotka-Volterra system

- Numerical tests of symmetry preservation: the rigid body

Perspectives

- Poisson integrators in solid mechanics (with L. Le Marrec and V. Carlier)

- Deformation theory of symplectic groupoids and plasma physics

Definition

M be a smooth manifold.

Recall that a Lie bracket on $\mathcal{C}^\infty(M)$ is a bilinear anti-symmetric map from $\mathcal{C}^\infty(M) \times \mathcal{C}^\infty(M) \rightarrow \mathcal{C}^\infty(M)$ such that

$$\{f, \{g, h\}\} + \{g, \{h, f\}\} + \{h, \{f, g\}\} = 0 \quad (\text{Jacobi relation}).$$

Definition (Flato, Lichnerowicz, Sternheimer, 1975)

A Poisson bracket on M is a Lie bracket on $\mathcal{C}^\infty(M)$ such that for all $f, g, h \in \mathcal{C}^\infty(M)$,

$$\{fg, h\} = f\{g, h\} + \{f, g\}h \quad (\text{Leibniz rule}).$$

Remark

Anti-symmetry of $\{.,.\} \Rightarrow$ for all $H \in \mathcal{C}^\infty(M)$, $\{H, H\} = 0$.

Consequence of the Leibniz rule: Poisson tensor of $\{\cdot, \cdot\}$

Lemma

An $\mathbb{R}$ -bilinear map $B: C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)$ arises from a smooth bivector field $\pi \in \Gamma(\bigwedge^2 TM)$ by the formula

$$B(f, g) = \langle \pi, df \wedge dg \rangle$$

if and only if B is skew-symmetric and verifies the Leibniz rule

$$B(fg, h) = fB(g, h) + B(f, g)h$$

Corollary

Any Poisson bracket $\{\cdot, \cdot\}$ gives rise to a smooth bivector field $\pi \in \Gamma(\bigwedge^2 TM)$. If $(x_i)_i$ are coordinates on M , $\pi_{i,j} = \{x_i, x_j\}$.

Remark

This holds because we assumed M to be finite dimensional.

Two consequences of the Jacobi relation

Proposition (Poisson brackets are particular cases of bivector fields)

There is a 1-to-1 correspondence between Poisson brackets on M and smooth bivector fields π such that the map

$$(f, g) \in \mathcal{C}^\infty(M)^2 \mapsto \langle \pi, df \wedge dg \rangle$$

satisfies the Jacobi relation.

Definition

Let $H \in \mathcal{C}^\infty(M)$. $f \in \mathcal{C}^\infty(M)$ is a first integral of H if $\{H, f\} = 0$.

Proposition (Poisson theorem)

$$\{H, f\} = \{H, g\} = 0 \Rightarrow \{H, \{f, g\}\} = 0.$$

Jacobi cyclic relation implies a Lie algebra structure on the first integrals of H .

A first example: symplectic manifolds

Let M be a symplectic manifold, meaning a manifold equipped with a closed 2-form ω such that for all $x \in M$, the map:

$$\omega^\flat: X \in T_x M \mapsto \omega_x(X, \cdot) \in T_x^* M$$

is a bijection, with inverse denoted by $\omega^\sharp$. The Poisson bracket of a symplectic manifold is

$$\{f, g\} = \langle df, \omega^\sharp(dg) \rangle.$$

Remark

Jacobi relation for this Poisson bracket is equivalent to the closedness of ω .

In Darboux coordinates (q, p) , the Poisson tensor is given by the matrix $\begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ and the corresponding Poisson bracket

$$\{f, g\}(q, p) = \frac{\partial f}{\partial q} \frac{\partial g}{\partial p} - \frac{\partial g}{\partial q} \frac{\partial f}{\partial p}.$$

Two more examples

2nd example: Lie-Poisson structure on $\mathfrak{g}^*$

Let $\mathfrak{g}$ be a Lie algebra. $M = \mathfrak{g}^*$ comes canonically equipped with the Poisson bracket

$$\forall f, g \in \mathfrak{g}^*, \quad \forall x \in \mathfrak{g}, \quad \{f, g\}(x) = \langle x, [d_x f, d_x g] \rangle$$

where $d_x f \in \mathfrak{g}^{**} \simeq \mathfrak{g}$.

3rd example: Cluster Poisson brackets

Let $A = (a_{ij})_{1 \leq i, j \leq n} \in \mathcal{M}_{n \times n}(\mathbb{R})$ anti-symmetric. Set $M = \mathbb{R}^n$ and

$$\forall x \in \mathbb{R}^n, \quad \forall 1 \leq i, j \leq n, \quad (\pi(x))_{ij} = a_{ij} x_i x_j.$$

Poisson morphisms

Definition

Let $(M_1, \{\cdot, \cdot\}_1)$ and $(M_2, \{\cdot, \cdot\}_2)$ two Poisson manifolds. A smooth map $\alpha: M_1 \rightarrow M_2$ is said to be a Poisson map, resp. an anti-Poisson map, if

$$\forall f, g \in \mathcal{C}^\infty(M_2), \quad \{\alpha^*f, \alpha^*g\}_1 = \alpha^*\{f, g\}_2$$

$$\text{resp. } \{\alpha^*f, \alpha^*g\}_1 = -\alpha^*\{f, g\}_2.$$

Example (Symplectic case)

Poisson automorphisms of a symplectic Poisson structure are called symplectomorphisms.

Example (Linear case)

Linear Poisson automorphisms of the dual of a Lie algebra are transpose maps of automorphisms of the Lie algebra.

Hamiltonian vector fields and flows

Definition (Hamiltonian vector field)

Identifying $\mathfrak{X}(M) \simeq \text{Der}(M)$, the Hamiltonian vector field of H is set to be

$$X_H: \begin{array}{ccc} \mathcal{C}^\infty(M) & \rightarrow & \mathcal{C}^\infty(M) \\ f & \mapsto & \pi(dH \wedge df) \end{array} .$$

Remark (Hamiltonian systems are local)

X_H at $x \in M$ only depend on the differential $d_x H$ of H at x .

Theorem

The Hamiltonian flow ϕ_t^H of H is a Poisson automorphism:

$$\forall f, g \in \mathcal{C}^\infty(M), \quad \forall t, \quad (\phi_t^H)^* \{f, g\} = \{(\phi_t^H)^* f, (\phi_t^H)^* g\}$$

Terminology

Hamiltonian flows are said to be inner automorphisms of π .

1st example of a Hamiltonian system: classical mechanics on $M = T^*X$

X	a smooth manifold	(the configuration space)
$\ \cdot\ $	a Riemannian metric on X	(the kinetic energy)
V	$\in \mathcal{C}^\infty(X)$	(the potential energy).

$$H: \begin{array}{ccc} T^*X & \rightarrow & \mathbb{R} \\ \zeta & \mapsto & \frac{1}{2m}\|\zeta\|^2 + V \circ \mathfrak{p}(\zeta) \end{array}$$

where: $\mathfrak{p}: T^*X \rightarrow V$ the cotangent projection
 $m > 0$ the mass.

Remark

The first integrals of H depend on the physics. For instance: angular momentum.

In cotangent coordinates (q, p) , the ODE of X_H is:

$$\begin{cases} \dot{q} = -\frac{p}{m} \\ \dot{p} = -\nabla V(q) \end{cases} \quad \text{i.e. 2nd law of Newton (1687) } m\ddot{q} = -\nabla V(q)$$

2nd example of a Hamiltonian system: mechanics on $\mathfrak{g}^*$ and the angular velocity of a rigid body

For the **canonical Poisson structure on $\mathfrak{g}^*$** , the Hamiltonian system associated to $H \in \mathcal{C}^\infty(\mathfrak{g}^*)$ is

$$\dot{x} = \text{ad}^*_{dH} x$$

In the case of $\mathfrak{so}(3) \simeq \mathbb{R}^3$, the Poisson bracket is

$$\{f, g\}(x) = (\nabla_x f)^T \cdot \begin{pmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{pmatrix} \cdot \nabla_x g$$

The Hamiltonian of the angular velocity of the rigid body is

$$H(x) = \frac{1}{2} \left(i_1(x_2^2 + x_3^2) + i_2(x_1^2 + x_3^2) + i_3(x_1^2 + x_2^2) \right)$$

$$\Rightarrow \dot{x} = -x \wedge Ix, \text{ where } I = \begin{pmatrix} i_1 & 0 & 0 \\ 0 & i_2 & 0 \\ 0 & 0 & i_3 \end{pmatrix} \text{ is the inertia tensor}$$

3rd example of a Hamiltonian system: Lotka-Volterra system on a cluster Poisson structure

the cluster Poisson bracket

$$\{f, g\}(x) = (\nabla_x f)^T \cdot \begin{pmatrix} 0 & x_1 x_2 & x_1 x_3 \\ -x_1 x_2 & 0 & x_2 x_3 \\ -x_1 x_3 & -x_2 x_3 & 0 \end{pmatrix} \cdot \nabla_x g$$

For the Hamiltonian

$$H(x) = \sum_{i=1}^3 x_i$$

being the sum of the species populations, the corresponding Hamiltonian system is

$$\Rightarrow \begin{cases} \dot{x}_1 = x_1(x_2 + x_3) \\ \dot{x}_2 = x_2(-x_1 + x_3) \\ \dot{x}_3 = -x_3(x_1 + x_2) \end{cases} \quad (1)$$

Remark (Vanhaecke et al., 2016)

The equation (1) has been proven to be part of a family of completely integrable systems.

First integrals

Theorem


Let $F \in \mathcal{C}^\infty(M)$ a first integral of H . The flow of H commutes with the one of F :

$$\forall s, t, \quad \phi_s^F \phi_t^H = \phi_t^H \phi_s^F$$

Corollary

The Hamiltonian is always a preserved quantity along a trajectory of its Hamiltonian flow.

In $\mathbb{R}^2$ equipped with $\pi = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, the data of the Hamiltonian levels of H is enough to compute any integral curve of the Hamiltonian vector field of H .



Hyperbolic pendulum
 $H(q, p) = \frac{p^2}{2} + \cos q$

Casimir functions

Definition

$C \in \mathcal{C}^\infty(M)$ is said to be a Casimir if it is a first integral for all smooth functions on M . Equivalently, its Hamiltonian vector field vanishes.

Example (The non-degenerate case)

A symplectic Poisson structure has no non-trivial Casimir.

Example (The linear case)

Any element of the center of a Lie algebra $\mathfrak{g}$ provides a Casimir on $\mathfrak{g}^*$.

Example (A quadratic case)

Let A be a $n \times n$ real anti-symmetric matrix and $(a_i)_{1 \leq i \leq n} \in \ker A$. Then, the function $x \in \mathbb{R}^n \mapsto \prod_{i=1}^n x_i^{a_i}$ is a Casimir of the cluster Poisson bracket associated to the matrix A .

Symplectic foliation

On a Poisson manifold M , not all the points can be connected by Hamiltonian flows. This foliates M .

Theorem

Any Poisson manifold (M, π) is foliated into symplectic leaves. This foliation is singular.

A foliation is in particular a partition of the manifold into immersed submanifolds.

Let $x \in M$ and $\mathcal{F}_x$ the leaf of x .

- ▶ $\mathcal{F}_x$ is a symplectic manifold.
- ▶ The immersion $\mathcal{F}_x \hookrightarrow M$ is a Poisson morphism.
- ▶ Let $y \in M$. The topology of $\mathcal{F}_y$ might differ a lot from the one of $\mathcal{F}_x$, even if x and y are close.

Remark

Any Casimir must be constant on a symplectic leaf.


Symplectic foliation of the examples

The symplectic Poisson structure

The symplectic foliation is made of one symplectic leaf: the whole space.

Lie-Poisson structure on $\mathfrak{g}^*$


It is given by coadjoint orbits.



A symplectic leaf of $\mathfrak{so}(3)^* \simeq \mathbb{R}^3$

Cluster Poisson brackets

Exercise !



A symplectic leaf of $\{x_i, x_j\} = x_i x_j$,

Properties of Hamiltonian flows

Let $H \in \mathcal{C}^\infty(M)$. Its Hamiltonian flow ϕ_t^H

- ▶ **preserves H :** $H \circ \phi_t^H = H$,
Hamiltonian flows are *conservative*.
- ▶ **preserves the Poisson tensor:** $\phi_t^H \star \pi = \pi$,
Hamiltonian flows are *inner automorphisms of the Poisson structure*.
- ▶ **stay on a leaf:** $\forall x \in M, \quad \phi_t^H(x) \in \mathcal{F}_x$,
this provides a *construction* of any symplectic leaf, cf control theory.

Symplectic Runge-Kutta


Define an integrator as a family: $\varphi_\epsilon: x_n \mapsto x_{n+1}$ depending on the time-step ϵ .

Definition

In the space $\{(q, p)\}$ of position-momenta, an integrator φ_ϵ is said to be **symplectic** if it is a Poisson morphism for the Poisson

structure $\pi(q, p) = \begin{pmatrix} 0 & -Id \\ Id & 0 \end{pmatrix} := J$.

Symplectic
integrators are used
in astrophysics,
molecular dynamics,
optics...



Virgo project, 2004

How to cook up a symplectic integrator ?

Here: $\pi = J$.

A Runge-Kutta integrator for $\dot{x} = J \cdot \nabla_x H$ is of the form


$$\begin{aligned}x_{n+1} &= x_n + \epsilon \sum_{i=1}^s b_i k_i \\ k_i &= J \cdot \nabla H(x_n + \epsilon \sum_{j=1}^s a_{ij} k_j)\end{aligned}$$

where $x := (q, p)$.


Theorem

*If : $\forall (i, j), b_i a_{ij} + b_j a_{ji} = b_i b_j$,
then the integrator is symplectic.*

Runge-Kutta Poisson ?



A Hamiltonian trajectory for a Poisson structure in $\mathbb{R}^3$



An integrator that preserves this Poisson structure

In the general case, it is not enough to preserve the Poisson structure. Such an integrator, even of high order, may provide bad numerical simulations: *it destroys the foliation*.

Exercises for next week

1. Prove that the cluster Poisson brackets are indeed Poisson brackets, namely, that the brackets verify the Jacobi relation.
2. Compute the symplectic foliation of $\mathbb{R}^3$ endowed with $\{x_1, x_2\} = x_1 x_2$, $\{x_1, x_3\} = 2x_1 x_3$, $\{x_2, x_3\} = x_2 x_3$. Namely: describe the partition of $\mathbb{R}^3$ induced by these Poisson brackets.

Bibliography for the first lecture

- ▶ *Symplectic Geometry and Analytical Mechanics*, Paulette Libermann and Charles-Michel Marke, Kluwer, 1987
→ Poisson geometry and mechanics
- ▶ Chapter 1 of *Poisson Structures and their Normal Forms*, Jean-Paul Dufour and Nguyen Tien Zung, Birkhäuser, 2005
→ Basics on Poisson geometry
- ▶ *Lectures on Poisson Geometry*, Marius Crainic, Rui Loja Fernandes, Ioan Mărcuț, American Mathematical Society, 2021
→ More involved Poisson geometry